



Wealth index and risk of childhood overweight and obesity: evidence from four prospective cohorts in Peru and Vietnam

Rodrigo M. Carrillo-Larco · J. Jaime Miranda · Antonio Bernabé-Ortiz

Received: 11 March 2015 / Revised: 3 November 2015 / Accepted: 12 November 2015
© Swiss School of Public Health (SSPH+) 2015

Abstract

Objectives To estimate the incidence and risk of childhood overweight and obesity according to socioeconomic status in Peruvian and Vietnamese school-aged children.

Methods Longitudinal data from the Young Lives study were analyzed. Exposure was wealth index in tertiles. Outcome was overweight and obesity. Cumulative incidence per 100 children-years, relative risks (RR), and 95 % confidence intervals (95 % CI) were calculated. A hierarchical approach, including child- and family-related variables, was followed to construct multivariable models.

Results The cumulative incidence of overweight and obesity was 4.8 (95 % CI 4.1–5.5) and 1.7 (95 % CI 1.3–2.2) in the younger and older Peruvian cohort, respectively; and in Vietnam 1.5 (95 % CI 1.2–1.8) and 0.3 (95 % CI 0.2–0.5), respectively. The incidence of overweight and obesity was higher at the top wealth index tertile in all samples. In the older cohorts, comparing highest versus bottom wealth index tertile, RR of overweight and obesity was four to nine times higher: 4.25 in

Peru (95 % CI 2.21–8.18) and 9.11 in Vietnam (95 % CI 1.07–77.42).

Conclusions The results provide important information for childhood obesity prevention in countries moving ahead with economic, epidemiological and nutritional transitions.

Keywords Peru · Vietnam · Developing countries · Pediatric obesity · Social class · Prospective studies

Introduction

Childhood obesity has become a public health issue worldwide (de Onis et al. 2010), and prevalence of overweight and obesity has increased about 47 % from 1980 to 2013 (Ng et al. 2014). When analyzed by socioeconomic indicators, the prevalence of childhood obesity has different patterns between developed and developing countries. Childhood obesity is higher among the wealthier in developing countries, yet it is more common among the poor in developed countries (Dinsa et al. 2012).

Several developing countries have gone through a rapid economic transition, and are now considered emerging economies; however, there are still numbers of people living under poverty. Overall, economic transition has a negative effect on the rates of childhood obesity (Cai 2014; Swinburn et al. 2011; Wang and Lim 2012). In terms of national averages of gross domestic product (GDP), Peru is classified as an upper-middle income country and Vietnam as lower-middle income country (The World Bank 2015). These socioeconomic features—economic transition and GDP—have affected health parameters of both countries. Peru undergoes epidemiological and nutritional transition (Chaparro and Estrada 2012; Huicho et al. 2009), with growing

Electronic supplementary material The online version of this article (doi:10.1007/s00038-015-0767-7) contains supplementary material, which is available to authorized users.

R. M. Carrillo-Larco · J. J. Miranda (✉) · A. Bernabé-Ortiz
CRONICAS Center of Excellence in Chronic Diseases,
Universidad Peruana Cayetano Heredia, Av. Armendáriz 497,
2do Piso, Miraflores, Lima 18, Peru
e-mail: jaime.miranda@upch.pe

J. J. Miranda
Department of Medicine, School of Medicine, Universidad
Peruana Cayetano Heredia, Lima, Peru

A. Bernabé-Ortiz
Escuela de Medicina, Universidad Peruana de Ciencias
Aplicadas - UPC, Lima, Peru

prevalence of childhood overweight and obesity whilst still hosting chronic malnutrition rates of 25 % (Alvarez-Dongo et al. 2012; Loret de Mola et al. 2014; Pajuelo-Ramirez et al. 2013). In Vietnam, the nutritional status of children has also changed: between 1992–1993 and 1997–1998, the prevalence of underweight decreased from 70 to 64 %; and prevalence of overweight and obesity is 20 % in school-aged children (Nguyen et al. 2013; Thang and Popkin 2003).

Childhood is a critical period to fight obesity as the effects of socioeconomic disparities on obesity risk appear and become stronger during childhood (Jansen et al. 2013). Moreover, the health consequences of childhood obesity at adulthood may be preventable (McMullen 2014). Overall, socioeconomic patterns are different between countries, and childhood is an important stage to prevent the effect of socioeconomic status on obesity. Therefore, studying children at different age ranges in countries where economic growth goes at different pace provides important scoping to identify key issues for the development of preventative strategies. This is particularly relevant as obesity prevention programs may have a different effect according to socioeconomic status (Beauchamp et al. 2014). Furthermore, previous studies have included surrogates to assess socioeconomic status such as parental educational attainment or household size (Wrotniak et al. 2012; Zadinska et al. 2012). Thus, there is need to explore the risk of childhood obesity according to socioeconomic status with a more comprehensive definition of socioeconomic status for developing countries, such as wealth index, which also has other strengths in comparison to other methods to assess socioeconomic status (Howe et al. 2012). Data to calculate wealth index are available for many countries as much of the work on this indicator has been developed with Demographic and Health Surveys. In addition, it is less susceptible to changes in income and expenditure (Howe et al. 2012).

In the present study, we contrast four prospective cohorts: two in Peru and two in Vietnam. The aim of the present study is to estimate the incidence of overweight and obesity according to socioeconomic status, and to assess whether socioeconomic status, as per wealth index, is a risk factor for the development of overweight and obesity in school-aged children in one lower-middle and one upper-middle income country, where risk factors for non-communicable diseases are on the rise.

Methods

Study design and setting

This is a secondary analysis of a longitudinal cohort: the Young Lives study, which is being conducted in four

developing countries: Ethiopia, India, Peru and Vietnam (Barnett et al. 2013).

The Young Lives study started in the year 2002 with two baseline groups in each country: a younger cohort included children aged 6–18 months and an older cohort included children aged 7–8 years. The follow-ups were conducted in the years 2006–2007 and 2009–2010 for both cohorts. Children in the younger cohort at the first and second follow-up were 4–5 and 7–8 years old, respectively; the corresponding figures for children in the older cohort are 11–12 and 14–15 years old.

Only information from Peru and Vietnam was included in this secondary analysis. The other countries were not included because preliminary analysis revealed low prevalence of overweight and obesity, preventing us from creating multivariable models. For example, in Ethiopia, in the younger cohort, 4.9 and 1.2 % of children were overweight or obese; in India, in the younger cohort, 1.4 % was overweight and 0.9 % obese. Moreover, we did not include the baseline round of the younger cohorts (children aged 6–18 months) because we targeted school-aged children. We included the first (hereafter known as *Baseline*) and second follow-ups of the younger cohorts, and the baseline assessment and second follow-up of the older cohorts.

Participants

Regarding this secondary analysis, the exclusion criteria for eligibility included missing values in key variables: child body mass index (BMI), age or sex, and household wealth index. Regarding the Peruvian younger cohort, 4.8 % of the participants were excluded at baseline; whilst with regard to the Peruvian older cohort, 0.6 % of the participants were excluded at baseline. The same exclusion criteria applied for Vietnam: at baseline 2.9 and 0.1 % were excluded in the younger and older cohort, respectively (Fig. 1).

Sampling and procedures

Exact details about sampling procedures and data collection methods have been published elsewhere (Barnett et al. 2013). To choose the participants in both cohorts, the Peruvian team followed a multi-stage, cluster-stratified, random sampling; whereas in Vietnam a multi-stage, purposive, random sampling was conducted (Online Resource 1) (The Young Lives Study 2008).

Variables

Outcome variable

The outcome was assessed at the second follow-up of either cohort. The outcome variable was the occurrence of




Fig. 1 Flowchart of the Peruvian and Vietnamese younger (2006–2007 to 2009–2010) and older (2002 to 2009–2010) cohorts. The Young Lives study. Further details about the attrition rate and follow-up of the Young Lives study have been published elsewhere

(Barnett et al. 2013). Baseline assessment of the younger cohort was conducted in 2006–2007 and of the older cohort it was conducted in 2002; follow-up of both cohorts was conducted in 2009–2010

overweight and obesity as per BMI, using age- and sex-specific cut-off points proposed by the World Obesity Federation (Cole and Lobstein 2012). We decided to combine overweight and obesity to have sufficient observations in both countries to fit multivariable models. Weight and height were measured in all children.

Exposure variable

The exposure variable was household wealth index in tertiles: bottom (poorest), middle and top (richest). The exposure was assessed at baseline of either cohort as was not expected to change markedly during follow-up. This wealth index was constructed as a simple average of three individual indexes: housing quality index (number of rooms, as well as floor, roof and wall material) consumer durables index (possession of radio, television, refrigerator, motorcycle, car, telephone, mobile phone and bike) and housing services index (source of drinking water, having electricity, type of toilet, and type of cooking fuel). A wealth index approach has become a common assessment method regarding socioeconomic status in developing countries and it is a useful measure of the household socioeconomic status (Howe et al. 2012). This variable was assessed with a face-to-face questionnaire answered by the household head or the mother.

Baseline assessment

Co-variables were assessed at baseline and with a face-to-face questionnaire answered by the household head or the mother. For the younger cohorts in Peru and Vietnam, we included: child age, child gender, birth weight (2500–4000 g, ≤ 2500 g, ≥ 4000 g), breastfeeding (Did you ever breastfeed the child?, yes/no), and number of meals the child had the previous day (< 5 , > 5 or 5), maternal nutritional status as per BMI [normal weight (≥ 18.5 and < 25), overweight (≥ 25 and < 30), obesity (≥ 30)], maternal education (none/primary, high school, higher education), and household location (rural/urban). Co-variables for the older cohorts in Peru and Vietnam included: child age, child gender, and birth order (first, second and third or more), maternal nutritional status as per BMI [normal weight (≥ 18.5 and < 25), overweight (≥ 25 and < 30), obesity (≥ 30)], maternal education (none/primary, high school, higher education), and household location (rural or urban).

Statistical analysis

To describe the exposure, outcome and co-variables, we used proportions and 95 % confidence intervals (95 % CI); means and standard deviations are also presented. Proportions were used to present categorical variables, while means were used

for numerical variables. Comparisons between categorical variables were performed using Chi-squared test. Cumulative incidence per 100 children-years and 95 % CI were calculated for developing overweight and obesity, after excluding those children who had overweight and obesity at baseline. We excluded these children because the incidence calculates the number of new cases over subjects at risk, thus subjects who already have the condition of interest should be excluded. Relative risk (RR) and 95 % CI were assessed using generalized lineal models with Poisson family and log link, fitting the model using robust standard errors to account for the cluster effect introduced by the sampling procedure. The analyses were conducted with STATA 11.0 (StataCorp, College Station, TX, USA).

To assure comparability between estimates resulting from younger or older cohorts, the multivariable models in both Peru and Vietnam were adjusted for the same co-variables. Four models were constructed in the multivariable analyses. The first, a crude model, only included the outcome and exposure variables. Model A added child's gender, age, and birth order. Given additional data availability in the younger cohorts, model A in this group also included birth weight, breastfeeding and number of meals the child had the previous day. Model B only included maternal education, maternal nutritional status, and household location. Model C included all previous variables. The cluster variable was included in the four models as a confounder. These models were fitted using a hierarchical approach, (Victora et al. 1997) conducted to understand whether the influence of socioeconomic status on the development of overweight and obesity is the same when controlling for child-related or family-related variables only.

Ethics

The Young Live Study had ethical approval from the ethics committee of Social Science division, University of Oxford; this approval was for the study as a whole. Further details on the ethics procedures have been published elsewhere (Young Lives, Ethics; Young Lives, The ethics of social research with children and families in young lives: Practical experiences).

Results

Sample description

The study included 4018 and 3856 children-years in the younger and older Peruvian cohort, respectively; and 5029 and 6787 children-years in the younger and older Vietnamese cohort, respectively (Fig. 1).

There were 1954 and 710 children at baseline in the younger and older Peruvian cohort, respectively. At baseline, mean age (\pm SD) of children in the younger cohort was 5.3 (\pm 0.4) years; 50.5 % were boys, and 69.5 % lived in urban settings. Mean age (\pm SD) of children in the older cohort, at baseline, was 7.9 (\pm 0.3) years, 54.1 % were boys, and they were predominately from urban settings (74.1 %).

In Vietnam, at baseline, there were 1944 and 999 children included in the younger and older cohort, respectively. Participants in the younger cohort had a mean age (\pm SD) of 5.3 (\pm 0.3) years, 51.6 % were boys, and the majority (79.5 %) lived in rural areas. At baseline, children in the older cohort had a mean age (\pm SD) of 8.0 (\pm 0.3) years, 50.2 % were boys, and 80.1 % lived in rural settings.

Nutritional status at baseline

At baseline in Peru, prevalence of overweight and obesity in the younger cohort was 16.2 % (95 % CI 14.5–17.9 %) and 5.4 % (95 % CI 4.0–6.4 %), respectively. There was more overweight and obesity at the top tertile of wealth index (28.3 %) compared to the bottom (18.4 %) and middle (17.8 %) tertiles ($p < 0.001$). The prevalence of childhood overweight and obesity in the older cohort was 15.4 % (95 % CI 12.7–18.1 %) and 3.3 % (95 % CI 2.0–4.7 %), respectively. There was more overweight and obesity as the wealth index grows up: 9.0, 22.0, and 25.4 % in the bottom, middle and top tertile, respectively ($p < 0.001$). Distribution of co-variables at baseline according to nutritional status and wealth index is presented in Online Resource 2 and 3, for both Peruvian cohorts.

At baseline in Vietnam, 5.8 % (95 % CI 4.6–7.1 %) were overweight and 4.7 % (95 % CI 3.6–5.9 %) were obese in the younger cohort. There was higher prevalence of overweight and obesity in the highest tertile of wealth index: 4.4, 4.8, and 21.4 % in the bottom, middle and top tertile, respectively ($p < 0.001$). Regarding the older cohort, 2.4 % (95 % CI 1.1–3.8 %) were overweight and 0.4 % (95 % CI 0.0–0.9 %) were obese. Also, there were more children with overweight and obesity at the top tertile of wealth index: 1.1 and 7.3 % in the bottom and top tertile, respectively ($p < 0.001$); there were none in the middle tertile. Distribution of co-variables at baseline according to nutritional status and wealth index is presented in Online Resource 4 and 5, for both Vietnamese cohorts.

Overweight and obesity incidence and comparisons by socioeconomic status

In Peru, the overall cumulative incidence of overweight and obesity was 4.8 (95 % CI 4.1–5.5) and 1.7 (95 % CI

Table 1 Cumulative incidence and 95 % confidence interval (95 % CI) of childhood overweight and obesity according to socioeconomic status at baseline

Variables	Younger cohort			Older cohort		
	WI: Bottom	WI: Middle	WI: Top	WI: Bottom	WI: Middle	WI: Top
Mother						
Maternal education						
None/primary	1.7 (1.1–2.8)	2.7 (1.7–4.2)	7.8 (4.7–12.9)	1.4 (0.9–2.4)	1.5 (0.7–2.9)	2.2 (0.7–6.7)
High school	1.9 (0.8–4.5)	4.7 (3.3–6.9)	8.8 (6.7–11.7)	1.2 (0.5–3.3)	1.3 (0.6–2.7)	2.7 (1.6–4.5)
Higher education	9.2 (2.3–36.8)	4.2 (2.1–8.3)	10.5 (7.8–14.2)	0.0 (0.0–0.0)	0.8 (0.1–5.4)	1.6 (0.7–3.3)
Maternal BMI						
Normal weight	1.8 (1.1–3.0)	3.1 (2.0–4.9)	7.9 (5.5–11.2)	1.2 (0.6–2.2)	1.3 (0.6–2.9)	1.3 (0.6–3.2)
Overweight	2.8 (1.5–5.0)	3.1 (2.0–4.9)	8.4 (6.2–11.5)	1.5 (0.8–3.1)	1.1 (0.5–2.4)	2.9 (1.8–4.8)
Obesity	1.2 (0.2–8.8)	8.1 (4.9–13.4)	11.9 (8.2–17.4)	1.0 (0.1–7.3)	2.0 (0.8–5.5)	2.3 (1.0–5.2)
Location						
Urban	2.1 (1.1–4.2)	4.3 (3.2–5.7)	9.2 (7.6–11.2)	1.8 (1.0–3.3)	1.2 (0.7–2.2)	2.3 (1.5–3.3)
Rural	2.0 (1.3–3.1)	2.2 (1.1–4.2)	7.0 (1.0–49.9)	1.1 (0.6–2.1)	1.5 (0.6–4.0)	0.0 (0.0–0.0)
Children						
Gender						
Male	2.5 (1.5–4.1)	3.3 (2.2–4.9)	9.4 (7.2–12.3)	0.8 (0.4–1.9)	1.0 (0.5–2.1)	2.3 (1.4–3.8)
Female	1.5 (0.9–2.9)	4.2 (2.9–6.0)	9.0 (6.9–11.8)	2.0 (1.2–3.3)	1.7 (0.9–3.2)	2.1 (1.2–3.8)
Age (years)^a						
4 years/6–7 years	2.5 (1.4–4.5)	3.3 (1.8–6.0)	4.6 (1.9–11.0)	2.0 (1.2–3.3)	1.0 (0.4–2.1)	2.4 (1.4–4.1)
5 years/8–10 years	1.8 (1.1–2.9)	3.8 (2.8–5.1)	9.5 (7.8–11.6)	0.8 (0.3–1.8)	1.6 (0.9–3.0)	2.1 (1.2–3.6)
6 years	0.0 (0.0–0.0)	5.9 (1.5–23.7)	12.9 (6.5–12.8)			
Birth weight (g)						
2501–3999	2.1 (1.3–3.3)	4.0 (3.0–5.3)	9.9 (8.2–12.1)	No data		
≤2500	1.8 (0.2–12.4)	2.5 (0.6–9.9)	5.8 (2.2–15.5)			
≥4000	0.0 (0.0–0.0)	4.9 (1.8–13.0)	6.7 (2.2–20.8)			
Breastfeeding						
Yes	2.0 (1.3–2.9)	3.8 (2.9–4.9)	9.2 (7.6–11.1)	No data		
No	20.0 (2.8–142.3)	0.0 (0.0–0.0)	5.8 (0.8–41.3)			
Child birth order						
First	No data			1.5 (0.6–3.6)	2.6 (1.4–4.8)	1.3 (0.6–2.9)
Second				0.7 (0.2–2.9)	0.0 (0.0–0.0)	2.5 (1.3–4.9)
Third or more				1.6 (0.9–2.7)	1.2 (0.5–2.6)	3.2 (1.8–5.7)
Number of meals the previous day						
5	1.8 (0.9–3.7)	5.3 (3.8–7.4)	8.5 (6.4–11.1)	No data		
<5	2.2 (1.3–3.7)	2.5 (1.4–4.4)	9.7 (6.3–14.9)			
>5	1.9 (0.8–4.5)	2.5 (1.3–4.7)	9.9 (7.1–13.8)			

The Young Lives study, younger (2006–2007 to 2009–2010) and older cohort (2002 to 2009–2010), Peru

^a Baseline age for either the younger or older cohort, respectively. WI = wealth index. Baseline assessment of the younger cohort was conducted in 2006–2007 and of the older cohort it was conducted in 2002; follow-up of both cohorts was conducted in 2009–2010

1.3–2.2) per 100 children-years for the younger and older cohort, respectively. The cumulative incidence of overweight and obesity was 9.2 (95 % CI 7.6–11.1) at the top wealth index tertile for the younger cohort, yet the incidence was smaller in the older cohort: 2.3 (95 % CI 1.6–3.3). Further details about the cumulative incidence according to co-variables and to wealth index are shown in Table 1.

In Vietnam, the overall incidence of overweight and obesity was 1.5 (95 % CI 1.2–1.8) per 100 children-years for the younger cohort and 0.3 (95 % CI 0.2–0.5) for the older cohort. The cumulative incidence of overweight and obesity was higher at the top wealth index tertile when compared to the middle and bottom tertile in either the younger or older cohort: 2.8 (95 % CI 2.0–3.8) and 0.7 (95 % CI 0.4–1.2), respectively. Further details about the

Table 2 Cumulative incidence and 95 % confidence interval (95 % CI) of childhood overweight and obesity according to socioeconomic status at baseline

Variables	Younger cohort			Older cohort		
	WI: Bottom	WI: Middle	WI: Top	WI: Bottom	WI: Middle	WI: Top
Mother						
Maternal education						
None/primary	0.8 (0.4–1.4)	1.4 (0.7–2.5)	1.0 (0.3–3.0)	0.2 (0.1–0.6)	0.3 (0.1–1.0)	0.6 (0.2–1.9)
High school	0.3 (0.0–2.0)	1.0 (0.5–1.9)	2.7 (1.7–4.2)	0.0 (0.0–0.0)	0.1 (0.0–0.6)	0.8 (0.4–1.6)
Higher education	0.0 (0.0–0.0)	3.0 (0.4–21.3)	5.5 (3.1–9.9)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	0.8 (0.2–3.3)
Maternal BMI						
Normal weight	0.7 (0.3–1.3)	1.2 (0.7–1.9)	2.7 (1.8–3.8)	0.0 (0.0–0.0)	0.2 (0.1–0.6)	0.4 (0.2–0.9)
Overweight	3.7 (1.2–11.5)	3.4 (1.1–10.5)	5.5 (2.6–11.5)	0.5 (0.1–3.5)	1.1 (0.2–7.7)	2.5 (1.1–5.6)
Obesity	0.0 (0.0–0.0)	8.9 (1.3–63.5)	11.8 (3.0–47.5)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	0.0 (0.0–0.0)
Location						
Urban	0.0 (0.0–0.0)	1.1 (0.4–3.5)	4.4 (2.9–6.6)	0.0 (0.0–0.0)	0.5 (0.1–3.9)	1.1 (0.6–1.9)
Rural	0.8 (0.4–1.3)	1.1 (0.7–1.8)	1.8 (1.1–3.0)	0.1 (0.0–0.4)	0.2 (0.1–0.5)	0.4 (0.1–1.0)
Children						
Gender						
Male	0.8 (0.4–1.7)	1.3 (0.7–2.2)	2.9 (1.9–4.5)	0.0 (0.0–0.0)	0.3 (0.1–0.8)	0.8 (0.4–1.6)
Female	0.7 (0.3–1.6)	0.9 (0.5–1.9)	2.6 (1.9–4.3)	0.3 (0.1–0.8)	0.2 (0.0–0.7)	0.6 (0.3–1.3)
Age (years)^a						
4 years/6–7 years	0.5 (0.1–2.0)	2.8 (1.4–5.5)	5.2 (2.7–10.0)	0.1 (0.0–0.6)	0.2 (0.0–0.7)	0.7 (0.4–1.6)
5 years/8 years	0.8 (0.5–1.5)	0.8 (0.4–1.4)	2.4 (1.7–3.4)	0.2 (0.0–0.7)	0.2 (0.1–0.8)	0.7(0.3–1.4)
6 years/–	0.0 (0.0–0.0)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	–	–	–
Birth weight (g)						
2501–3999	0.9 (0.5–1.7)	1.2 (0.7–1.9)	2.7 (1.9–3.7)	No data		
≤2500	1.4 (0.2–9.8)	0.0 (0.0–0.0)	3.6 (1.2–11.1)			
≥4000	2.3 (0.3–16.0)	1.7 (0.2–12.2)	3.6 (0.9–14.5)			
Breastfeeding						
Yes	0.8 (0.4–1.3)	1.1 (0.7–1.7)	2.8 (2.1–3.8)	No data		
No	0.0 (0.0–0.0)	0.0 (0.0–0.0)	0.0 (0.0–0.0)			
Child birth order						
First	No data			0.2 (0.1–0.8)	0.3 (0.1–1.0)	0.7 (0.3–1.6)
Second				0.2 (0.0–1.3)	0.4 (0.1–1.2)	0.9 (0.4–1.9)
Third or more				0.0 (0.0–0.0)	0.0 (0.0–0.0)	0.5 (0.2–1.6)
Number of meals the previous day						
5	0.7 (0.2–2.1)	1.1 (0.5–2.4)	1.5 (0.7–3.2)	No data		
<5	0.8 (0.4–1.6)	0.6 (0.3–1.5)	0.8 (0.3–2.5)			
>5	0.6 (0.1–3.9)	2.1 (1.0–4.2)	4.7 (3.3–6.8)			

The Young Lives study, younger (2006–2007 to 2009–2010) and older cohort (2002 to 2009–2010), Vietnam

^a Baseline age for either the younger or older cohort, respectively. WI = wealth index. Baseline assessment of the younger cohort was conducted in 2006–2007 and of the older cohort it was conducted in 2002; follow-up of both cohorts was conducted in 2009–2010

cumulative incidence by wealth index tertiles according to co-variables are shown in Table 2.

Risk of overweight and obesity according to socioeconomic status

The bottom wealth index tertile was considered as the reference group in multivariable analyses. In both younger

cohorts, from Peru and Vietnam, there was no evidence of a higher risk of overweight and obesity by socioeconomic status. On the contrary, there was higher overweight and obesity risk in both—Peru and Vietnam—older cohorts. Peruvian children in the older cohort showed higher risks of overweight and obesity: twofold on middle vs. bottom tertile and fourfold on the top vs. bottom tertile of wealth index (Fig. 2). Corresponding RR estimates for Vietnam




Fig. 2 Relative risk and 95 % confidence interval (95 % CI) of childhood overweight and obesity. The Young Lives study, younger (2006–2007 to 2009–2010) and older (2002 to 2009–2010) Cohorts, Peru and Vietnam. Baseline assessment of the younger cohort was conducted in 2006–2007 and of the older cohort it was conducted in 2002; follow-up of both cohorts was conducted in 2009–2010. Younger cohort adjusted for child gender, child age at baseline, birth

weight, breastfeeding, number of meals the previous day at baseline, maternal education at baseline, maternal nutritional status at baseline and location at baseline. Older cohort adjusted for child gender, child age at baseline, child birth order, maternal education, maternal nutritional status at baseline and location at baseline. *Y* younger cohort, *O* older cohort

were 7.69 (middle vs. bottom tertile of wealth index) and 9.11 (top vs. bottom tertile of wealth index) (Fig. 2). Further details about the multivariable models are shown in Table 3 and Online Resource 6.

Risk of overweight and obesity according to socioeconomic status: a hierarchical approach

In Peru, the magnitude of the association between child-related variables and overweight and obesity tended to be higher than maternal ones, except for the older cohort when comparing top versus bottom tertile (Table 3); however, in Vietnam, the magnitude of the association between child-related variables and overweight and obesity was higher than maternal ones in the older cohort when comparing top versus bottom tertile (Online Resource 6).

In the Peruvian younger cohort, the risk of overweight and obesity was statistically significant when adjusting only for child-related variables; with regard to the Peruvian older cohort, the risk of overweight and obesity was significant when adjusting for either child- or family-related variables (Table 3). In the Vietnamese younger cohort, the risk of overweight and obesity was statistically significant when adjusting only for child-related variables and when comparing bottom versus top tertile; in the Vietnamese older cohort, when comparing bottom versus top tertile, there was higher risk of overweight and obesity after

adjusting for either child-related or family-related variables (Online Resource 6).

Discussion

Main findings

Our results show that, relative to lowest socioeconomic strata, the incidence of childhood overweight and obesity is higher among the wealthier group. Overall, there is a dissimilar impact of socioeconomic status on childhood overweight and obesity in two developing countries in different world regions; this is supported by the different cumulative incidences found in Peru when compared to Vietnam, also by the magnitude of the relative risks reported. The effect of socioeconomic status on the risk of developing overweight and obesity appears to be different depending on the child age, being much stronger among older children. In addition, it appears that maternal characteristics plays a more important role among younger than older children.

The results—higher risk of obesity among wealthier individuals—could also contribute to picture how economic transition affects the risk of childhood obesity: in the older cohort, the risk of overweight and obesity was much higher in Vietnam than in Peru among individuals at the

Table 3 Relative risk and 95 % confidence interval (95 % CI) of childhood overweight and obesity

		Crude	Model A	Model B	Model C
Younger cohort					
Bottom	<i>n</i> = 1695	1	1	1	1
Middle		1.65 (1.14–2.38)	1.6 (1.07–2.25)	0.98 (0.65–1.49)	0.97 (0.64–1.47)
Bottom	<i>n</i> = 1695	1	1	1	1
Top		3.75 (2.69–5.23)	3.36 (1.34–4.81)	1.36 (0.87–2.14)	1.31 (0.83–2.06)
Older cohort					
Bottom	<i>n</i> = 769	1	1	1	1
Middle		2.68 (1.56–4.62)	2.95 (1.71–5.10)	2.23 (1.33–3.73)	2.37 (1.42–3.93)
Bottom	<i>n</i> = 749	1	1	1	1
Top		3.52 (2.09–5.94)	3.84 (2.26–6.52)	4.21 (2.19–8.09)	4.25 (2.21–8.18)

The Young Lives study, younger (2006–2007 to 2009–2010) and older (2002 to 2009–2010) cohort, Peru

Baseline assessment of the younger cohort was conducted in 2006–2007 and of the older cohort it was conducted in 2002; follow-up of both cohorts was conducted in 2009–2010. Younger cohort: model A included child gender, child age at baseline, birth weight, breastfeeding, number of meals the previous day at baseline; model B included only maternal education at baseline, maternal nutritional status at baseline and location (rural/urban) at baseline; model C included all the previous variables. Older cohort: model A included child gender, child age at baseline and child birth order; model B only included maternal education (brought from round 2), maternal nutritional status at baseline and location (rural/urban) at baseline; model C included all the previous variables. In bold, $p < 0.05$

top wealth index tertile. As Vietnam continues its path towards upper-middle-income country, the current Peruvian status, its risk estimates for overweight and obesity will likely decrease and get closer to those observed in Peru. Similarly, if Peru keeps its economic growth towards a high-income country, the higher risk observed among the better-off relative to the worse-off in terms of socioeconomic indicators might disappear.

Previous studies

The overall cumulative incidence of childhood obesity assessed from kindergarten to eighth grade (1998–2007) in the United States was 11.9 per 100 children-years, and opposite to our results, the smaller incidence was observed among the top socioeconomic quintile (Cunningham et al. 2014). In Europe, greater deprivation means a higher prevalence of childhood overweight (Knai et al. 2012; Stamatakis et al. 2010) and a similar pattern has been described in Australia (O’Dea et al. 2014). However, results from other developing countries are similar to ours (Friedman et al. 2009; Hyska et al. 2014; Navti et al. 2014; Schooling et al. 2010; Wrotniak et al. 2012; Zadzinska et al. 2012). Thus, the effect of socioeconomic status on childhood obesity is different between developed and developing countries; and within the latter there are different patterns of childhood obesity according to socioeconomic status.

Evidence from transitional countries in Europe is valuable. A cross-sectional study in Albania reported there were more children with overweight and obesity in urban settings, versus rural areas. Although they aimed to

assess the demographic correlates of childhood overweight and obesity, they failed to include a more comprehensive parameter of socioeconomic status relying only in location (Hyska et al. 2014). In Ukraine, children whose mother was from middle/upper social class have 60 % higher odds of overweight and obesity (Friedman et al. 2009). A study in Poland that included cohorts of children followed through different settings of economic and political transition revealed similar results: the risk of overweight and obesity in boys was 34 % higher for those whose mothers had secondary and post-secondary education, versus elementary or basic vocation plus higher (Zadzinska et al. 2012).

African countries have also provided evidence to support the different effect of socioeconomic status on childhood obesity. Botswana is undergoing a rapid nutritional transition, and Wrotniak et al. in a cross-sectional study reported similar results to ours, though with smaller estimates: children in private school had 300 % higher odds of being obese versus their peers in public schools; also, children in families with high assets had 260 % higher odds, versus children living in families with low assets (Wrotniak et al. 2012). In a cross-sectional study in Cameroon, Navti et al. found that children at the highest socioeconomic status had higher odds of being overweight (Navti et al. 2014); their estimates were similar to what we found in Vietnam, yet they used a different method to assess socioeconomic status (Navti et al. 2014).

These studies provide evidence that childhood obesity is more common among the wealthier groups regardless of where the developing country is, and despite the definition of socioeconomic status.

Results' interpretation

The impact of socioeconomic status on overweight and obesity are independent of maternal characteristics among the older children in both Peru and Vietnam but not in the younger ones. This observation is supported by our results, when considering regression model B comparing bottom versus top tertile, the results are significant for the older cohorts, but not for the younger cohorts. However, which maternal characteristic exerts a stronger effect on childhood obesity needs to be further addressed. In an attempt to explore which maternal characteristic plays a more important role in the risk of childhood overweight and obesity according to socioeconomic status, we conducted post hoc analyses with the Peruvian younger cohort. Comparing bottom with middle wealth index tertile, when only adding to the crude model maternal education, the relative risk went from 1.65 ($p = 0.007$) to 1.30 ($p = 0.191$); yet when only including maternal nutritional status, the estimate increased to 1.69 ($p < 0.001$). When running the same post hoc analysis by comparing bottom versus top tertile, the new estimates were: 2.24 ($p < 0.001$) and 3.18 ($p < 0.001$). Thus, it seems that at younger ages, maternal education exerts a major impact on the risk of childhood overweight and obesity according to socioeconomic status. Maternal education has not been much exploited in previous intervention programs (Hillier-Brown et al. 2014).

In this study, we found sufficient evidence that the incidence of overweight and obesity is higher in the top wealth index category, and it is much stronger among older children. Therefore, our findings call for the implementation of relevant and appropriate obesity prevention strategies alongside improvements in socioeconomic status, e.g., poverty reduction strategies, the so-called lifting people out poverty, should anticipate mechanisms for not pushing people into obesity. These prevention strategies call for active and creative thinking and meaningful conversations across fields, for example, between those concerned with health and development in the international strategies agenda. At the in-country level, our research findings call for interactions between ministries of economy, welfare, education, health and social protection, among others, to convey and implement such preventative strategies.

We provide evidence that the burden of childhood overweight and obesity in two developing countries is much higher among the rich, in comparison to the poor; similar results have been reported for cardiovascular risk factors, including obesity, in India (Subramanian et al. 2013). However, it would be inappropriate to fight childhood overweight and obesity only in these high-risk populations, because health needs go beyond just

socioeconomic status (Busingye et al. 2014; Razak and Subramanian 2014). Therefore, as people progress towards wealthier socioeconomic status, interventions and prevention strategies must be creative to avoid social development coming along with growing rates of childhood overweight and obesity.

Strengths and limitations

This study benefits of a prospective design, which rules out reverse causality. We have included a wide sample size of children in two developing countries, at different development stage, and in different world regions. In addition, we have included several potential confounders, which were also the same when comparing the younger or older cohorts. Moreover, we used a comprehensive method to assess socioeconomic status—wealth index—as previous reports used characteristics such as household size and parental education (Zadzinska et al. 2012), or type school and assets (Wrotniak et al. 2012). The wealth index approach is commonly used in developing countries and it has some strengths: it is less prone to change according to seasonality patterns of consumption, and it does assess the socioeconomic status at the household level (Howe et al. 2012).

However, limitations must be addressed too. First, due to data availability, we did not include the same variables across the younger and older cohorts. Thus, we were not able to compare children within a country. Post hoc analysis showed that, when including in the final model only variables shared in common by the four cohorts, results were quite identical (data not shown). Second, the sampling in either country excluded the richest settings. This is particularly relevant for Vietnam as these households were poorer than the average Vietnamese household. Thus, the highest socioeconomic status presented in this study does not necessarily represent the highest socioeconomic status in the general population. If the richest group had been included, our estimates could have been greater. Thus, our results are conservative and alert of a possible higher risk of overweight and obesity among the better-off. Third, the sampling procedures were not exactly the same in Peru and Vietnam. The Vietnamese team followed a purposive procedure at some stages of the sampling; though they still cover the diversity of Vietnamese children. Fourth, the loss of respondents through study rounds could have added attrition bias. Nonetheless, it has been reported that there is low attrition bias and that the attrition rate has a minor impact on any result (Barnett et al. 2013). Finally, we did not include an important confounder such as physical activity as this variable was not available at baseline. In post hoc analysis when we included physical activity assessed at follow-up, the models did not vary much (data

not shown). We believe this—lack of adjustment for physical activity—did not affect our results and we are still able to hypothesize that maternal characteristics play a more important role in obesity prevention among younger children.

Conclusions

There is strong evidence of higher risk of overweight and obesity in older children in Peru and Vietnam, especially among those at the highest wealth index; however, no strong association was found among younger children. The effect of socioeconomic status on childhood overweight and obesity was stronger in Vietnam than in Peru. These results are important in the current global childhood obesity epidemic, and in particular for those countries undergoing economic, epidemiological and nutritional transition.

Acknowledgments The data used in this publication come from Young Lives, a 15-year study of the changing nature of childhood poverty in Ethiopia, India (Andhra Pradesh), Peru and Vietnam (<http://www.younglives.org.uk>). Young Lives is funded by UK aid from the Department for International Development (DFID) and co-funded from 2010 to 2014 by the Netherlands Ministry of Foreign Affairs. The views expressed here are those of the author(s). They are not necessarily those of Young Lives, the University of Oxford, DFID or other funders.

The authors are thankful to Daniel Lopez de Romaña Forga, PhD for his comments on earlier versions of this manuscript.

Compliance with ethical standards

Funding RMC-L, AB-O, JJM are part of the CRONICAS Center of Excellence in Chronic Diseases, supported by the National Heart, Lung, and Blood Institute Global Health Initiative under the contract Global Health Activities in Developing Countries to Combat Non-Communicable Chronic Diseases (Project Number 268200900033C-1-0-1). AB-O is supported by a Wellcome Trust Research Training Fellowship in Public Health and Tropical Medicine (Grant Number: 103994/Z/14/Z).

Conflict of interest The authors declare no conflict of interests.

References

- Alvarez-Dongo D, Sanchez-Abanto J, Gomez-Guizado G et al (2012) Overweight and obesity: prevalence and determining social factors of overweight in the Peruvian population (2009–2010). *Revista peruana de medicina experimental y salud publica* 29(3):303–313
- Barnett I, Ariana P, Petrou S et al (2013) Cohort profile: the Young Lives study. *Int J Epidemiol* 42(3):701–708. doi:10.1093/ije/dys082
- Beauchamp A, Backholer K, Magliano D et al (2014) The effect of obesity prevention interventions according to socioeconomic position: a systematic review. *Obes Rev: Off J Int Assoc Study* 15(7):541–554. doi:10.1111/obr.12161
- Busingye D, Arabshahi S, Subasinghe AK et al (2014) Do the socioeconomic and hypertension gradients in rural populations of low- and middle-income countries differ by geographical region? A systematic review and meta-analysis. *Int J Epidemiol* 43(5):1563–1577. doi:10.1093/ije/dyu112
- Cai W (2014) Nutritional challenges for children in societies in transition. *Curr Opin Clin Nutr Metab Care* 17(3):278–284. doi:10.1097/mco.0000000000000042
- Chaparro MP, Estrada L (2012) Mapping the nutrition transition in Peru: evidence for decentralized nutrition policies. *Revista panamericana de salud publica = Pan Am J Public Health* 32(3):241–244
- Cole TJ, Lobstein T (2012) Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes* 7(4):284–294. doi:10.1111/j.2047-6310.2012.00064.x
- Cunningham SA, Kramer MR, Narayan KM (2014) Incidence of childhood obesity in the United States. *N Engl J Med* 370(17):1660–1661. doi:10.1056/NEJMc1402397
- de Onis M, Blossner M, Borghi E (2010) Global prevalence and trends of overweight and obesity among preschool children. *Am J Clin Nutr* 92(5):1257–1264. doi:10.3945/ajcn.2010.29786
- Dinsa GD, Goryakin Y, Fumagalli E et al (2012) Obesity and socioeconomic status in developing countries: a systematic review. *Obes Rev: Off J Int Assoc Study* 13(11):1067–1079. doi:10.1111/j.1467-789X.2012.01017.x
- Friedman LS, Lukyanova EM, Serdiuk A et al (2009) Social-environmental factors associated with elevated body mass index in a Ukrainian cohort of children. *Int J Pediatr Obes: IJPO: Off J Int Assoc Study* 4(2):81–90. doi:10.1080/17477160802449264
- Hillier-Brown FC, Bamba CL, Cairns JM et al (2014) A systematic review of the effectiveness of individual, community and societal level interventions at reducing socioeconomic inequalities in obesity amongst children. *BMC Public Health* 14(1):834. doi:10.1186/1471-2458-14-834
- Howe LD, Galobardes B, Matijasevich A et al (2012) Measuring socio-economic position for epidemiological studies in low- and middle-income countries: a methods of measurement in epidemiology paper. *Int J Epidemiol* 41(3):871–886. doi:10.1093/ije/dys037
- Huicho L, Trelles M, Gonzales F et al (2009) Mortality profiles in a country facing epidemiological transition: an analysis of registered data. *BMC Public Health* 9:47. doi:10.1186/1471-2458-9-47
- Hyska J, Mersini E, Mone I et al (2014) Prevalence and demographic correlates of overweight and obesity among children in a transitional Southeastern European population. *J Community Health*. doi:10.1007/s10900-014-9888-9
- Jansen PW, Mensah FK, Nicholson JM et al (2013) Family and neighbourhood socioeconomic inequalities in childhood trajectories of BMI and overweight: longitudinal study of Australian children. *PLoS One* 8(7):e69676. doi:10.1371/journal.pone.0069676
- Knai C, Lobstein T, Darmon N et al (2012) Socioeconomic patterning of childhood overweight status in Europe. *Int J Environ Res Public Health* 9(4):1472–1489. doi:10.3390/ijerph9041472
- Loret de Mola C, Quispe R, Valle GA et al (2014) Nutritional transition in children under five years and women of reproductive age: a 15-years trend analysis in Peru. *PLoS One* 9(3):e92550. doi:10.1371/journal.pone.0092550
- McMullen S (2014) Childhood obesity: the impact on long-term risk of metabolic and CVD is not necessarily inevitable. *Proc Nutr Soc* 73(3):389–396. doi:10.1017/s0029665114000111
- Navti LK, Ferrari U, Tange E et al (2014) Contribution of socioeconomic status, stature and birth weight to obesity in Sub-Saharan Africa: cross-sectional data from primary school-age children in Cameroon. *BMC Public Health* 14:320. doi:10.1186/1471-2458-14-320

- Ng M, Fleming T, Robinson M et al (2014) Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. doi:10.1016/s0140-6736(14)60460-8
- Nguyen PV, Hong TK, Hoang T et al (2013) High prevalence of overweight among adolescents in Ho Chi Minh City, Vietnam. *BMC Public Health* 13:141. doi:10.1186/1471-2458-13-141
- O’Dea JA, Chiang H, Peralta LR (2014) Socioeconomic patterns of overweight, obesity but not thinness persist from childhood to adolescence in a 6-year longitudinal cohort of Australian schoolchildren from 2007 to 2012. *BMC Public Health* 14:222. doi:10.1186/1471-2458-14-222
- Pajuelo-Ramirez J, Sanchez-Abanto J, Alvarez-Dongo D et al (2013) Overweight, obesity and chronic mal nutrition in 6 to 9 year-old children in Peru, 2009–2010. *Revista peruana de medicina experimental y salud publica* 30(4):583–589
- Razak F, Subramanian S (2014) Commentary: socioeconomic status and hypertension in low- and middle-income countries: can we learn anything from existing studies? *Int J Epidemiol* 43(5):1577–1581. doi:10.1093/ije/dyu159
- Schooling CM, Yau C, Cowling BJ et al (2010) Socio-economic disparities of childhood body mass index in a newly developed population: evidence from Hong Kong’s ‘Children of 1997’ birth cohort. *Arch Dis Child* 95(6):437–443. doi:10.1136/adc.2009.168542
- Stamatakis E, Wardle J, Cole TJ (2010) Childhood obesity and overweight prevalence trends in England: evidence for growing socioeconomic disparities. *Int J Obes* (2005) 34(1):41–47. doi:10.1038/ijo.2009.217
- Subramanian SV, Corsi DJ, Subramanyam MA et al (2013) Jumping the gun: the problematic discourse on socioeconomic status and cardiovascular health in India. *Int J Epidemiol* 42(5):1410–1426. doi:10.1093/ije/dyt017
- Swinburn BA, Sacks G, Hall KD et al (2011) The global obesity pandemic: shaped by global drivers and local environments. *Lancet* 378(9793):804–814. doi:10.1016/s0140-6736(11)60813-1
- Thang NM, Popkin B (2003) Child malnutrition in Vietnam and its transition in an era of economic growth. *J Hum Nutr Diet: Off J Br Diet Assoc* 16(4):233–244
- The World Bank (2015) Data by country. <http://data.worldbank.org/country>. Accessed 18 Nov 2015
- Victora CG, Huttly SR, Fuchs SC et al (1997) The role of conceptual frameworks in epidemiological analysis: a hierarchical approach. *Int J Epidemiol* 26(1):224–227
- Wang Y, Lim H (2012) The global childhood obesity epidemic and the association between socio-economic status and childhood obesity. *Int Rev Psychiatry (Abingdon, Engl)* 24(3):176–188. doi:10.3109/09540261.2012.688195
- Wrotniak BH, Malet L, Maruapula SD et al (2012) Association between socioeconomic status indicators and obesity in adolescent students in Botswana, an African country in rapid nutrition transition. *Pediatr Obes* 7(2):e9–e13. doi:10.1111/j.2047-6310.2011.00023.x
- The Young Lives Study (2008) An assessment of the Young Lives sampling approach in Peru. <http://www.younglives.org.uk/files/technical-notes/an-assessment-of-the-young-lives-sampling-approach-in-peru>. Accessed 18 Nov 2015
- Young Lives. An assessment of the Young Lives sampling approach in Vietnam. <http://www.younglives.org.uk/files/technical-notes/an-assessment-of-the-young-lives-sampling-approach-in-vietnam>
- Young Lives. Young Lives methods guide sampling. <http://www.younglives.org.uk/files/methods-guide/methods-guide-sampling>
- Zadzinska E, Rosset I, Koziel S et al (2012) Frequency of under- and overweight among children and adolescents during the economic transition in Poland. *Homo: internationale Zeitschrift für die vergleichende Forschung am Menschen* 63(3):216–232. doi:10.1016/j.jchb.2012.03.001